

Molina Healthcare of Washington Medicaid Private Duty Nursing Prior Authorization Request Form Phone Number: (800) 869-7175

Fax Number: (800) 767-7188

MEMBER INFORMATION										
Plan:	Molina Medicaid (If Molina is secondary, please include a copy of the denial from primary insurance)									
Member Name:							DOB:	/	/	
Member ID#:							Phone:	()) –	
Service Type:	Elec ⁻	tive/Rou	itine 🗆 Expe	edited/U	rgent					
REFERRAL/SERVICE TYPE REQUESTED										
Diagnosis Code & Description:										
CPT/HCPC Code & Description:										
CPT/HCPC Code & Description:										
90 DOS SPAN ONLY For continuation requests, the start date is always the day after the last authorization ends						DS From: / / to / /				
PROVIDER INFORMATION										
Requesting Provide	r Name:				NPI#:			TIN#		
Servicing Provider or	Facility:				NPI#:			TIN#		
Contact at Req Provider's								1	l	
Phone N	umber:	ımber: () –				Fax mber: () -				
CLINICAL DOCUMENTATION TO SUPPORT NEED FOR PRIVATE DUTY NURSING (PDN)										
Signed and dated physician order for PDN [Please submit: Home Health Certification and Plan of Ca of Health and Human Services, HCFA Form: OBM 0938-0				· · ·				□ Submitted		
Current history and physical (recent hospital admissions/discharge summ Current treatment plan and treatment re Current nursing care plan - Most recent r Recent daily nursing notes Emergency medical plan				s) s □ S (two weeks) □ S □ S		Submitted Submitted Submitted Submitted Submitted		 Not Submitted Not Submitted Not Submitted Not Submitted Not Submitted 		
90 DAY SUMMARY/including changes □ Submitted □ Not Submitted										
If YES, please desc	ribe:									

CLINICAL PRESENTATION (check all that apply)					
Frequency of assessments (to include vital signs, interventions to support patient care, health					
status assessment, etc.) :					
Once per eight hour shift					
□ 2-3 times per eight hour shift					
□ Hourly or more often					
Behavioral health, cognition, developmental monitoring:					
□ Non-verbal, infrequent speech, or difficult to understand					
□ Self-abusive behavior, risk of self-harm, and intervention required					
□ Sleep disturbance and patient awake more than three hours per night					
Combative, confused, or disoriented behavior that impacts self-management; patient obese					
□ Combative, confused, or disoriented behavior that impacts self-management					
Respiratory:					
☐ More than eight hours per day					
□ Less than eight hours per day					
□ Nebulizer therapy					
□ More frequent than every four hours					
Every 4-24 hours					
□ Less frequent than daily, but at least once every seven days					
□ Chest Physiotherapy – percussion, high-frequency chest wall oscillation vest, cough assist					
device, etc. □ More than once per hour					
Every 1-4 hours					
 Less than every four hours, but at least daily 					
□ Oxygen management					
□ Oxygen humidification, tracheal, no ventilator					
Oxygen needed at least weekly, based on pulse oximetry					
□ Suctioning					
□ Tracheal suctioning at least once every two hours					
Tracheal suctioning daily, but less than every two hours					
□ Nasal or oral suctioning daily					
Tracheostomy management					
□ Tracheostomy management with complications (skin breakdown, replacement needed)					
Tracheostomy management, no complications					
🗆 Ventilator management					
Continuous ventilator use					
Ventilator use for 12 or more hours per day					
Ventilator use for 7-12 hours per day					
Ventilator use for less than seven hours per day					
Interventions in place for active weaning					
Ventilator weaning achieved; requires ongoing post-weaning monitoring and management					
\square Ventilator on standby, respiratory assistance, or used at night for less than one hour					

Skilled Nursing Needs:						
🗆 Blood draw						
	Peripheral line					
	Less than twice per week					
□ Infusion therapy						
□ Blood or blood product						
Chemotherapy infusion						
Central line access and management						
Pain medication infusion						
□ Intravenous Infusion (IV antibiotics, etc.), including	infusion administration and monitoring					
for infusion reactions						
□ Infusions more than every four hours						
□ Infusions less than every four hours						
□ Non-infusion medication						
□ Insulin administration						
□ Non-insulin medication injectable administ						
Medication administration at least every tw	o nours, requiring clinical monitoring					
□ Activity of Daily Living (ADL)/Therapy support						
BedboundWh	eeichdir user Ambulatory					
□ Total/partial lift, weight 55-125 pounds	undo					
□ Total/partial lift, weight greater than 125 pounds						
□ ADL support needed more than four hours per day to maximize patient's independence						
Body cast management Cast or brace management						
 Splinting management, including removal ar 	ad raplacement, at least every eight hours					
□ Communication deficit; nurse to support the						
□ Range of motion exercises at least every eig						
 Range of motion exercises at least every eight hours Physical therapy program at least three hours per day; occupational therapy program 						
at least four hours per day	rs per day, occupational therapy program					
□ Nutrition management						
☐ Enteral nutrition with complications, requires administration of feeding, residual check,						
adjustment or placement of tube, and assessment or management of complications						
Enteral nutrition without complications						
□ Gastrostomy tube care, uncomplicated						
□ Nasogastric tube care, uncomplicated						
Partial parenteral nutrition with central line care						
☐ Total parenteral nutrition with central line co						
□ Skin and wound care management						
□ Burn care						
Ostomy care, at least once per day						
Postsurgical care, within 45 days of surgery						
□ Stage one or two wound management, at le	east once per day					
□ Stage three or four wound management, at least once per day						
□ Stage three or four wound management at least once per day, and multiple wound sites						
Prescribed topical medication application a						
U Wound vacuum management						

□ Seizure control that requires nursing intervention/management

- □ Seizures lasting less than three minutes, at least four times per week
 - □ Seizures lasting 3-5 minutes, at least four times per week
 - □ Seizures lasting 3-5 minutes, one to four times per day
 - □ Seizures lasting 3-5 minutes, more than five times per day
 - □ Seizures lasting more than five minutes, or clustered seizures, or seizure activity without regaining consciousness, at least four times per week
 - Seizures lasting more than five minutes, or clustered seizures, or seizure activity without regaining consciousness, one time or more per day, requiring rectal medication
 - □ Seizures lasting more than five minutes, or clustered seizures, or seizure activity without regaining consciousness, one time or more per day, requiring IM or IV medication

ADDITIONAL INFORMATION

List: